

Data Source:
Shoreline Type
GeoBC Coastal Resource Shorezone Database, 2008
Base Information
1:20,000 GeoBC Terrain Resource Information
Management (TRIM) Database

1:20,000

0 0.25 0.5 1

Kilometers



Shoreline Habitat

The Habitat Type provides a simplified picture of the "look" of the unit overall, based on the detailed biophysical attributes that have been mapped. The Habitat Type category is a summary of the observations of both the unit's biological and geomorphological features.

Each Habitat Type has a definition that includes the typical substrate, wave exposure and biobands. For example, for the

Each Habitat Type has a definition that includes the typical substrate, wave exposure and biobands. For example, for the Semi-exposed, Immobile substrate Habitat Type, part of the definition of that class is a certain combination of the most likely biobands and indicator species present at a bedrock shoreline with no mobile sediment present.

Each Habitat Type has typical biological features (including both an indicator species list and typical associated biobands). To determine the Habitat Type, the biomapper looks at the along-shore Units as designated and described by the physical mapper, and

1. records the observations of the biobands in the unit and looks for indicator species,
2. assigns a bio-(wave) exposure category,
3. writes the observed measured information and

- 3.□reviews the physical mapped information, and
- 4.□assigns the Habitat Type that best describes the unit.

The Habitat Type is based on the whole unit and is similar to the physical mappers use of the 'Coastal Class' category, in that the detailed across-shore data are summarized into one attribute. The simplified category describes the features of the whole unit.

Habitat Type is a summary of the biophysical classification of the whole shore unit, based on:

Habitat Type is a summary of the biophysical classification of the whole shore unit, based on:

- the biobands observed,
- the wave exposure as indicated by the bands, and

- the wave exposure as indicated by the bands, and
- the substrate types in the unit.

Legend Definitions
CC - Coastal Classification number

Wave Exposure
© 2014 Pearson Education, Inc.

E - Exposed - Very high wave exposure, open ocean swellsm usually fetches >500km
VE - Very Exposed - Extreme high wave exposure
SE - Semi-Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km

SE - Semi Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km
P - Protected - Low wave exposure, sheltered inlets, usually fetches less than 10km
SP - Semi Protected - Moderate wave exposure, partly sheltered, usually fetches 10-50km

SP - Semi Protected - Moderate wave exposure, partly sheltered, usually fetches 10-50km
VP - Very Protected - Very low wave exposure, fetches < 1km, sheltered anchorages at heads of bays and inlets

| Table WCVI. GOES WITH BIO AREAS WCVI, SCVI, WCVINorth, JdF Habitat Classification for “Exposure Bio” (EXP_BIO) and “Habitat Observed” (HAB_OBS) based on visible macro-biota assemblages for the West Coast Vancouver Island bio-mapping. | | | | | | | | | |
|--|--------------------------------|------------------------------|--|---|---|---|---|---------------------------------|---------------------|
| SUBSTRATE | BEDROCK/BOULDER | BEDROCK/BOULDER | BEDROCK/BOULDER | BEDROCK/BOULDER | SAND & GRAVEL | SAND & GRAVEL | SAND/MUD | SEDIMENT | BEDROCK OR SEDIMENT |
| OASTAL LASSES | 1-20 | 1-20 | 1-23, 32, 33 | 1-23, 33 | 24, 25, 26, 32 | 24, 25, 26, 32 | 27, 28, 29, 30, 31 | 24 - 30 | |
| POSURE XP. BIO) | E | SE | SP | P, VP | SP | P, VP | SP, P, VP | SE, E | VP, P, SP |
| HABITAT SERVED AB_OBS) | 2 | 3 * | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| upper | <i>Verrucaria</i> | <i>Verrucaria</i> | <i>Verrucaria</i> | <i>Verrucaria</i> | <i>Verrucaria</i> | <i>Verrucaria</i> | marsh grasses & rushes | | |
| | | <i>Enteromorpha</i> | <i>Enteromorpha</i> | <i>Enteromorpha</i> | <i>Enteromorpha</i> | <i>Enteromorpha</i> | | | |
| | | | | | | | <i>Saficornia virginica</i> | | |
| | <i>Balanus glandula</i> | <i>Balanus glandula</i> | <i>Balanus glandula</i> | <i>Balanus glandula</i> | <i>Balanus glandula</i> | <i>Balanus glandula</i> | <i>Balanus glandula</i> | | |
| | <i>Peltvetiopsis limitata</i> | <i>Fucus distichus</i> | <i>Fucus distichus</i> | <i>Fucus distichus</i> | <i>Fucus distichus</i> | <i>Fucus distichus</i> | <i>Fucus distichus</i> | | |
| middle | <i>Semibalanus cariosus</i> | <i>Semibalanus cariosus</i> | <i>Semibalanus cariosus</i> | | <i>Semibalanus cariosus</i> | | | | |
| | | | | | | | | | |
| | <i>Pollicipes polymerus</i> | | <i>Mytilus trossulus</i> | <i>Mytilus trossulus</i> | <i>Mytilus trossulus</i> | <i>Mytilus trossulus</i> | <i>Mytilus trossulus</i> | | |
| | | | <i>Ulva/ Ulvaria spp.</i> | <i>Ulva/ Ulvaria spp.</i> | <i>Ulva/ Ulvaria spp.</i> | <i>Ulva/ Ulvaria spp.</i> | <i>Ulva/ Ulvaria spp.</i> | | |
| mid/low | <i>Mytilus californianus</i> | <i>Mytilus californianus</i> | | | | | | | |
| | | | <i>Microcladia/Iridaea type mixed reds</i> | <i>Gigartina/Odonthalia type mixed reds</i> | | |
| | <i>Postelsia palmaeformis</i> | | | | | | | | |
| | | | <i>Hedophyllum sessile</i> | | | | | | |
| | | | <i>Codium fragile</i> | <i>Codium fragile</i> | | <i>Codium fragile</i> | | | |
| lower | <i>Lessoniopsis littoralis</i> | | | <i>Laminaria saccharina</i> | <i>Laminaria saccharina</i> | <i>Laminaria saccharina</i> | <i>Laminaria saccharina</i> | | |
| | | | | <i>Egregia menziesii</i> | | | | | |
| | <i>Laminaria setchellii</i> | | | | | | | | |
| | | | | <i>Laminaria groenlandica</i> | <i>Laminaria groenlandica</i> | | <i>Laminaria groenlandica</i> | | |
| | <i>Alaria nana</i> | | | <i>Alaria marginata</i> | | | <i>Alaria marginata</i> | | |
| | | | | <i>Eisenia arborea</i> | | | | | |
| | <i>Lithothamnion</i> | | | <i>Lithothamnion</i> | | | <i>Lithothamnion</i> | | |
| | | | | <i>Sargassum muticum</i> | | | <i>Sargassum muticum</i> | | |
| | | | | <i>Agarum sp</i> | <i>Agarum sp</i> | <i>Agarum sp</i> | <i>Agarum sp</i> | | |
| | | | | <i>Phyllospadix soudertii</i> | | | | | |
| subtidal | | | | <i>Macrocystis integrifolia</i> | <i>Macrocystis integrifolia</i> | <i>Macrocystis integrifolia</i> | <i>Macrocystis integrifolia</i> | <i>Macrocystis integrifolia</i> | |
| | | | | <i>Nereocystis luetkeana</i> | <i>Nereocystis luetkeana</i> | | <i>Nereocystis luetkeana</i> | | |
| | | | | <i>Strongylocentrotus</i> | <i>Strongylocentrotus</i> | | <i>Strongylocentrotus</i> | | |
| | | | | <i>composita</i> | <i>composita</i> | | <i>composita</i> | | |

