



Shoreline Habitat

The Habitat Type provides a simplified picture of the "look" of the unit overall, based on the detailed biophysical attributes that have been mapped. The Habitat Type category is a summary of the observations of both the unit's biological and geomorphological features.

Each Habitat Type has a definition that includes the typical substrate, wave exposure and biobands. For example, for the Semi-exposed, Immobile substrate Habitat Type, part of the definition of that class is a certain combination of the most likely biobands and indicator species present at a bedrock shoreline with no mobile sediment present.

How is Habitat Type determined?

Each Habitat Type has typical biological features (including both an indicator species list and typical associated biobands). To determine the Habitat Type, the biomapper looks at the along-shore Units as designated and described by the physical mapper, and 1. reviews the typical biological features of the biobands in the unit and looks for indicator species,

2. assigns a bio-stratigraphic exposure category,

3. reviews the physical mapped information, and

4. assigns the Habitat Type that best describes the unit.

The Habitat Type is based on the whole unit and is similar to the physical mappers use of the 'Coastal Class' category, in that the detailed across-shore data are summarized into one attribute. The simplified category describes the features of the whole unit.

Habitat Type is a summary of the biophysical classification of the whole shore unit, based on:

- the substrate types in the unit,
- the wave exposure as indicated by the bands, and
- the substrate types in the unit.

Legend Definitions

CC - Coastal Classification number

Wave Exposure

E - Exposed - High wave exposure, open ocean swellism usually fetches >500km

VE - Very Exposed - Extreme high wave exposure

SE - Semi Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km

P - Protected - Low wave exposure, sheltered inlets, usually fetches less than 10km

SP - Semi Protected - Moderate wave exposure, partly sheltered, usually fetches 10-50km

VP - Very Protected - Very low wave exposure, fetches < 1km, sheltered anchorages at heads of bays and inlets

Table WCVL GOES WITH BIO_AREAS WCVL, SCVL, WCVNORTH, JdF
Habitat Classification for "Exposure Bio" (EXP_BIO) and "Habitat Observed" (HAB_OBS) based on visible macro-bio assemblages for the West Coast
Vancouver Island bio-mapping.

MAJOR SUBSTRATE	BEDROCK/BOULDER	BEDROCK/BOULDER	BEDROCK/BOULDER	BEDROCK/BOULDER	SAND & GRAVEL	SAND & GRAVEL	SAND/MUD	SEDIMENT	BEDROCK OR SEDIMENT
COASTAL CLASSES	1-20	1-20	1-23, 32, 33	1-23, 33	24, 25, 26, 32	24, 25, 26, 32	27, 28, 29, 30, 31	24 - 30	
EXPOSURE (EXPOSED)	E	SE	SP	P, VP	SP	P, VP	SP, P, VP	SE, E	VP, P, SP
HABITAT OBSERVED (HAB_OBS)	2	3 *	4	5	6	7	8	9	10
upper	Terranea	Yerminius	Yerminius	Yerminius	Yerminius	Yerminius	Yerminius	Yerminius	Yerminius
	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha	Enteromorpha
middle									
lower									

