

| Vancouver Island bio-mapping. | | | |
|----------------------------------|-------------------------|---------------------------------------|----|
| MAJOR SUBSTRATE | BEDROCK/BOULDER | BEDROCK/BOULDER | |
| COASTAL CLASSES | 1-20 | 1-20 | |
| EXPOSURE (EXP_BIO) | Е | SE | |
| HABITAT OBSERVED (HAB_OBS) | 2 | 3 * | |
| upper | Verrucaria | Verrucaria | |
| | | Enteromorpha | ŀ |
| | Balanus glandula | Balanus glandula | |
| | Pelvetiopsis limitata | Fucus distichus | |
| middle | 1 crrenopono riminara | 1 HERD CHOROTHIS | f |
| | Semibalanus carriosus | Semibalanus carriosus | ļ |
| | Pollicipes polymerus | | j |
| | | | 1 |
| mid/low | Mytilus californianus | Mytilus californianus | |
| | | Microcladia/Iridea type mixed reds | t |
| | Postelsia palmaeformis | | |
| | | Hedophyllum sessile | |
| | | Codium fragile | |
| lower | Lessoniopsis littoralis | | |
| | | Egregia menziesii | |
| | Laminaria setchellii | Laminaria setchellii | |
| | | Laminaria groenlandica | |
| | Alaria nana. | Alaria marginata. | 4 |
| | | Eisenia arborea | |
| | Lithothamnion | Lithothamnion | |
| | | | |
| | | Agarum sp | 4 |
| | | Phyllospadix scouleri | |
| subtidal | | Macrocystis integrifolia | Ì |
| | Nereocystis luetkeana | Nereocystis luetkeana | j |
| | | Strongylocentrotus franciscanus | : |
| | | | ť |
| | 1 | • | 14 |
| | | | |