Data Source: Shoreline Type GeoBC Coastal Resource Shorezone Database, 2008 Base Information 1:20,000 GeoBC Terrain Resource Information Management (TRIM) Database 1:20,000 0 0.25 0.5 1 Kilometers		
Legend		
• Unit Break Points	Mobile/Partially Mobile Substrates	
Undefined	6 - Sand & Gravel - CC 24-26, 32 -SP	
mmobile Substrates	7 - Sand & Gravel - CC 24-26,32 - VP/P	The Habitat Type
🔨 1 - Bedrock - CC 1-20 - VE	8 - Estuary or Sand/Mud - CC 27-31 - VP/P/SP	been mapped. T features.
🔨 2 - Bedrock - CC 1-20 - E	✓ 9 - Sediment - CC 21 - 30 - SE/E	Each Habitat Typ Semi-exposed, Ir
🔨 3 - Bedrock/Boulder - CC 1-23, 32, 33 - S	SE Current Dominated	biobands and ind
4 - Bedrock/Gravel - CC 1-23, 33 - SP	10 - Bedrock or Sediment - CC 34 - VP/P/SP	How is Habitat Ty

• 4 - Bedrock/Gravel - CC 1-23, 33 - SH 5 - Bedrock/Gravel - CC 1-23,33 - P/VP Tidal Lagoon 11 - Bedrock or Sediment - CC 35 - VP/P/SP 
 CC
 Type
 CC
 Type

 Rock Shore Types - characterized by a lack of clastic sediments such as gravel or sand.
 Sediment Shore Types - have substrates that have little or no bedcrock cropping out

 1
 Lock Ramp, Wide
 21 Cravel Elst, Wide

 Rock Shore Types - characterized by a lack of clastic sediments such as grave

 1
 Rock Ramp, Wide

 2
 Rock Platform Wide

 3
 Rock Cliff Narrow

 4
 Rock Ramp, Narrow

 5
 Rock Platform Narrow

 Rock and Sediment Shore Types - rock and pockets of clastic sediments

 6
 Ramp with Gravel Beach Wide

 21
 Gravel Flat, Wide

 22
 Gravel Flat, Wide

 23
 Gravel Beach

 23
 Gravel Flat or Fan

 24
 Sand and Gravel Flat or Fan, Wide

 25
 Sand and Gravel Beach


 26
 Sand and Gravel Beach
26 Sand and Gravel Flat or Fan, Narrow 27 Sand Beach, Wide 28 Sand Flat Rock and Sediment Shore Types - rock and pockets of clastic6Ramp with Gravel Beach, Wide7Platform with Gravel Beach, Wide8Cliff with Gravel Beach9Ramp with Gravel Beach, Narrow10Platform with Gravel Beach, Narrow11Ramp with Sand and Gravel Beach, Wide12Platform with Sand and Gravel Beach, Wide13Cliff with Sand and Gravel Beach, Wide14Ramp with Sand and Gravel Beach, Narrow15Platform with Sand and Gravel Beach, Narrow16Ramp with Sand Beach, Wide17Platform with Sand Beach, Wide18Cliff with Sand Beach19Ramp with Sand Beach, Narrow20Platform with Sand Beach, Narrow 29 Mud Flat
 30 Sand Beach, Narrow
 31 Estuaries
 Man-Made Materials
 32 Man-made, permeable
 33 Man-made, impermeable
 Current Dominated
 34 Channel 34 Channel 35 Tidal Lagoon

103A.027

Mould Rock

Ρ

• The biobands observed, • the wave exposure as indicated by the bands, and • □ the substrate types in the unit.



SUBSTRATE STABILITY MAJOR SUBSTRATE	IMMOBILE SUBSTRATES			MOBILE OR PARTIALLY MOBILE SUBSTRATES				CURRENT- DOMI- NATED	TIDAL IAGOON	
	BEDROCK	BEDROCK/BOULDER	BEDROCK/GRAVEL	BEDROCK/GRAVEL	SAND & GRAVEL	SAND & GRAVEL	SAND/MUD	SEDIMENT	BEDROCK OR SEDIMENT	BEDROCK OR SEDIMENT
COASTAL CLASSES	1-20	1-23, 32, 33	1-23, 33	1-23, 33	24 – 30, 32 no SAL band	24 – 30, 32 no SAL band	24 - 30, 31 has SAL band	24-30	34	35
EXPOSURE (EXP BIO)	E	SE	SP	VP, P	SP	VP, P	VP, P, SP	SE, E	VP, P, SP	VP, P, SP
COMMUNITY CODE (HAB OBS)	2	3	4	5	6	7	8	9	10	11
upper	Verrucaria Balanus glandula	Verrucaria Enteromorpha Balanus glandula	Verrucaria Enteromorpha Balanus glandula	Verrucaria Enteromorpha Balanus giandula	Verrucaria Enteromorpha Balanus glandula	Verrucaria Enteromorpha Balanus glandula	grasses & rushes Salicornia virginica Balanus glandula			Balanus glandul
	, i i i i i i i i i i i i i i i i i i i	Fucus distichus	Fucus distichus	Fucus distichus	Fucus distichus	Fucus distichus	Fucus distichus	no visible	tidal current	Fucus distichus
middle	Pollicipes polymerus Mytilus californianus Semibalanus carriosus	Mytilus californianus Semibalanus carriosus	Mytilus trossulus* Semibalanus carriosus	Mytilus trossulus *	Semibalanus carriosus		Mytilus trossulus**	macrobiota due to sediment mobility	dominated; may be a Protected wave exposure but shows an	ponded water in
mid/low			Ulva/ Ulvaria spp.	Ulva/ Ulvaria spp.	Ulva/ Ulvaria spp.	Ulva/ Ulvaria spp.	Ulva/ Ulvaria	1	assemblage of indicator species	lagoon creates narrow intertidal
	Alaria 'nana' morph	Hedophyllum sessile Phyllospadix scouleri							from higher wave exposures. Assemblage	and a reduced biota in brackish water, may have
lower	Lessoniopsis littoralis Lithothamnion	Alaria 'marginata' morph Lithothannion	Laminaria groenlandica <b>Laminaria saccharina</b> Alaria 'marginata' morph Lithothamnion	Laminaria saccharina	Laminaria groenlandica Laminaria saccharina Alaria 'marginata' morph Lithothamnion	Laminaria saccharina			observed is 'anomalous' for the wave energy of the site.	dominated at outflow
subtidal	Nereocystis luetkeana	Nereocystis luetkeana Macrocystis integrifolla Agarum spp. Strongylocentrotus franciscanus	Nereocystis luetkeana Macrocystis integrifolia Agarum spp. Strongylocentrotus franciscanus Zostera marina	Macrocystis integrifolia Agarum spp. Zostera marina	Nereocystis luetkeana Macrocystis integrifolia Agarum spp. Strongylocentrotus franciscanus Zostera marina	Macrocystis integrifolia Agarum spp. <b>Zostera marina</b>	Zostera marina			

