

9 - Sediment - CC 21 - 30 - SE/E 2 - Bedrock - CC 1-20 - E 3 - Bedrock/Boulder - CC 1-23, 32, 33 - SE Current Dominated

10 - Bedrock or Sediment - CC 34 - VP/P/SP 4 - Bedrock/Gravel - CC 1-23, 33 - SP

5 - Bedrock/Gravel - CC 1-23,33 - P/VP

CC	Туре		cc	Туре
Rock Shore Types - characterized by a lack of clastic sediments such as gravel or sand.			Sediment 9	Shore Types - have substrates that have little or no bedcrock cropping out
	1 Rock Ramp, Wide		21	Gravel Flat, Wide
	2 Rock Platform Wide		22	Gravel Beach
	3 Rock Cliff Narrow		23	Gravel Flat or Fan
	4 Rock Ramp, Narrow		24	Sand and Gravel Flat or Fan, Wide
	5 Rock Platform Narrow		25	Sand and Gravel Beach
Rock a	nd Sediment Shore Types - rock and pockets of clastic sediments		26	Sand and Gravel Flat or Fan, Narrow
	6 Ramp with Gravel Beach, Wide		27	Sand Beach, Wide
	7 Platform with Gravel Beach, Wide		28	Sand Flat
	8 Cliff with Gravel Beach		29	Mud Flat
	9 Ramp with Gravel Beach, Narrow		30	Sand Beach, Narrow
	10 Platform with Gravel Beach, Narrow		31	Estuaries
	11 Ramp with Sand and Gravel Beach, Wide		Man-Made Materials	
	12 Platform with Sand and Gravel Beach, Wide		32	Man-made, permeable
	13 Cliff with Sand and Gravel Beach		33	Man-made, impermeable
	14 Ramp with Sand and Gravel Beach, Narrow		Current Dominated	
	15 Platform with Sand and Gravel Beach, Narrow		34	Channel
	16 Ramp with Sand Beach, Wide		35	Tidal Lagoon
	17 Platform with Sand Beach, Wide			
	18 Cliff with Sand Beach			
	19 Ramp with Sand Beach, Narrow			
	20 Platform with Sand Beach, Narrow			

Semi-exposed, Immobile substrate Habitat Type, part of the definition of that class is a certain combination of the most likely biobands and indictor species present at a bedrock shoreline with no mobile sediment present.

How is Habitat Type determined? Each Habitat Type has typical biological features (including both an indicator species list and typical associated biobands).

To determine the Habitat Type, the biomapper looks at the along-shore Units as designated and described by the physical mapper, and 1. □ records the observations of the biobands in the unit and looks for indicator species,

2. □ assigns a bio-(wave) exposure category,
3. □ reviews the physical mapped information, and
4. □ assigns the Habitat Type that best describes the unit.

The Habitat Type is based on the whole unit and is similar to the physical mappers use of the 'Coastal Class' category, in that the detailed across-shore data are summarized into one attribute. The simplified category describes the features of the whole unit.

Habitat Type is a summary of the biophysical classification of the whole shore unit, based on: • □ the biobands observed,

• the wave exposure as indicated by the bands, and • □ the substrate types in the unit.

Legend Definitions CC - Coastal Classification number

E - Exposed - Very high wave exposure, open ocean swellsm usually fetches >500km VE - Very Exposed - Extreme high wave exposure SE - Semi Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km P - Protected - Low wave expsoure, sheltered inlets, usually fetches less than 10km SP - Semi Protected - Moderate wave expsoure, partly sheltered, usually fetches 10-50km

species assemblages are as diagnostic as species' presence. Community Code type 1 (VE - very exposed) occurs only on the southwest coast of Moresby Island. VP - Very Protected - Very low wave exposure, fethces < 1km, sheltered anchorages at heads of bays and inletes

EXPOSURE (EXP BIO) COMMUNITY CODE (HAB_OBS) VP, P, SP SE, E VP, P, SP VP, P 8 | 9 | 10 grasses & rushes Salicornia Enteromorpha Enteromorpha Enteromorpha Enteromorpha Enteromorpha **virginica** Balanus glandula Fucus distichus tidal current dominated; may Balanus glandula Fucus distichus Balanus glandula middle be a protected wave exposure but shows an assemblage of indicator species Politcipes polymerus Mytilus trossulus Mytilus trossulus Mytilus trossulus Semibalanus carriosus Semibalanus carriosus [Semibalanus carriosus] Semibalanus carriosus Semibalanus carriosus Ulva/ Ulvaria spp. Ulva/ Ulvaria spp. Ulva/Ulvaria spp. Ulva/ Ulvaria spp. intertidal from higher macrobiota wave exposures. Hedophyllum sessile due to sediment mobility Alaria 'nana' morph] Alaria 'nana' morph Assemblage observed is Codium fragile Codium fragile Codium fragile Phyllospadix scouleri 'anomalous' for Egregia menziesii the wave energy Laminaria setchelli minaria setchelli] Laminaria setchelli Laminaria groenlandica Laminaria saccharina Laminaria saccharina Laminaria groenlandica Laminaria groenlandica lush foliose coralline diverse mixed red algae Laminaria saccharina reds: Bosstella/ Calliarthron/Corallina Alaria 'marginata'morph Alaria 'marginata'morph Alaria 'marginata'morph reocystis luetkeana Nereocysti s luetkeana Nereocystis luetkeana lereocystis luetkeana Macrocystis integrifolia Macrocystis integrifolia Macrocystis integrifolia Macrocystis integrifolia Agarum spp.Agarum spp. Agarum spp. Agarum spp. Strongylocentrotus franciscanus Strongylocentrotus Strongylocentrotus franciscanus franciscanus Zostera marina Zostera marina * Bolding indicates diagnostic species used to distinguish "communities". Square brackets [] enclose species at VE AB_OBS 1 which may be present but are in reduced abundance and size. Note that the absence of

