



Legend

Unit Break Points

Undefined

Immobile Substrates

- 1 - Bedrock - CC 1-20 - VE
- 2 - Bedrock - CC 1-20 - E
- 3 - Bedrock/Boulder - CC 1-23, 32, 33 - SE
- 4 - Bedrock/Gravel - CC 1-23, 33 - SP
- 5 - Bedrock/Gravel - CC 1-23, 33 - PNP

Mobile/Partially Mobile Substrates

- 6 - Sand & Gravel - CC 24-26, 32 - SP
- 7 - Sand & Gravel - CC 24-26, 32 - VP/P
- 8 - Estuary or Sand/Mud - CC 27-31 - VP/P/SP
- 9 - Sediment - CC 21 - 30 - SE/E
- 10 - Bedrock or Sediment - CC 34 - VP/P/SP
- 11 - Bedrock or Sediment - CC 35 - VP/P/SP

Current Dominated

Tidal Lagoon

CC - Coastal Classification number

CC Type	CC Type
Rock Shores - Rides characterized by a lack of classic sediments such as gravel or sand.	Sediment Shores - Types have substrates that have little or no bedrock cropping out.
1) Bank Ramp, Wide	21) Gravel Flat, Wide
2) Rock Platform, Wide	22) Gravel Beach
3) Bank Cliff, Narrow	23) Gravel Flat or Fan
4) Bank Ramp, Narrow	24) Sand and Gravel Flat or Fan, Wide
5) Bank Platform, Narrow	25) Sand and Gravel Beach
6) Bank with Gravel Beach, Wide	26) Sand and Gravel Flat or Fan, Narrow
7) Platform with Gravel Beach, Wide	27) Sand Beach, Wide
8) Cliff with Gravel Beach, Wide	28) Mud Flat
9) Beach with Gravel Beach, Narrow	29) Mud Flat
10) Platform with Gravel Beach, Narrow	30) Estuary
11) Beach with Sand and Gravel Beach, Wide	31) Sand Beach, Narrow
12) Cliff with Sand and Gravel Beach, Wide	32) Estuary
13) Beach with Sand and Gravel Beach, Narrow	33) Man-made, permeable
14) Platform with Sand and Gravel Beach, Narrow	34) Charcoal
15) Beach with Sand Beach, Wide	35) Tidal Lagoon
16) Platform with Sand Beach, Wide	
17) Cliff with Sand Beach	
18) Beach with Sand Beach, Narrow	
19) Platform with Sand Beach, Narrow	

Shoreline Habitat

The Habitat Type provides a simplified picture of the "look" of the unit overall, based on the detailed biophysical attributes that have been mapped. The Habitat Type category is a summary of the units biological and geomorphological features. Each Habitat Type has a definition that includes the typical substrate, wave exposure and biobands. For example, for the Semi-exposed, Immobile substrate Habitat Type, part of the definition of that class is a certain combination of the most likely biobands and indicator species present at a bedrock shoreline with no mobile sediment present.

How is Habitat Type determined?

Each Habitat Type has typical biological features (including both an indicator species list and typical associated biobands). To determine the Habitat Type, the biomapper looks at the along-shore Units as designated and described by the physical mapper, and

1. records the observations of the biobands in the unit and looks for indicator species,
2. assigns a bio-wave exposure category,
3. reviews the physical mapped information, and
4. assigns the Habitat Type that best describes the unit.

The Habitat Type is based on the whole unit and is similar to the physical mappers use of the 'Coastal Class' category, in that the detailed cross-shore data are summarized into one attribute. The simplified category describes the features of the whole unit.

Habitat Type is a summary of the biophysical classification of the whole shore unit, based on:

- the biobands observed,
- the wave exposure as indicated by the bands, and
- the substrate types in the unit.

Legend Definitions

CC - Coastal Classification number

Wave Exposure

- E - Exposed - Very high wave exposure, open ocean swells usually fetches >500km
- VE - Very Exposed - Extreme high wave exposure
- SE - Semi Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km
- P - Protected - Low wave exposure, sheltered inlets, usually fetches less than 10km
- SP - Semi Protected - Moderate wave exposure, partly sheltered, usually fetches 10-50km
- VP - Very Protected - Very low wave exposure, fetches < 1km, sheltered anchorages at heads of bays and inlets

Table MIDCOAST and NORTH COAST project area which includes BCO AREAS CC, IS and NC. The Species/wave exposure/ substrate table for Habitat Classification (HAB, OBS), for the Mid-coast BC study area, from Johnstone Strait/Central Coast Mapping Regions 5, 6 and 7.

SUBSTRATE STABILITY MAJOR SUBSTRATE	IMMOBILE SUBSTRATES					MOBILE OR PARTIALLY MOBILE SUBSTRATES			CURRENT-DOMINATED	TIDAL LAGOON
	BEFROCK	BEFROCK/BOULDER	BEFROCK-GRAVEL	BEFROCK-GRAVEL	SAND & GRAVEL	SAND & GRAVEL	SAND/SED.	SEDIMENT		
COASTAL EXPOSURE	1-20	1-23, 32, 33	1-23, 33	1-23, 33	24-30, 32 no SAL band	24-30, 32 no SAL band	24-30, 31 has SAL band	24-30	34	35
COMMUNITY CODE (HAB OBS)	E	3	4	5	6	7	8	9	10	11
upper	<i>Fernaria</i>	<i>Fernaria</i> <i>Enteromorpha</i>	<i>Fernaria</i> <i>Enteromorpha</i>	<i>Fernaria</i> <i>Enteromorpha</i>	<i>Fernaria</i> <i>Enteromorpha</i>	<i>Fernaria</i> <i>Enteromorpha</i>	<i>grasses & rushes</i> <i>Sargassum</i> <i>Ulva</i>			
middle	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Semibalanus cariosus</i>	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Semibalanus cariosus</i>	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Altilia prolifera</i> <i>Ulva</i> <i>Libinia</i> spp.	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Altilia prolifera</i> <i>Ulva</i> <i>Libinia</i> spp.	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Altilia prolifera</i> <i>Ulva</i> <i>Libinia</i> spp.	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Altilia prolifera</i> <i>Ulva</i> <i>Libinia</i> spp.	<i>Balanus glandula</i> <i>Mutinus californicus</i> <i>Altilia prolifera</i> <i>Ulva</i> <i>Libinia</i> spp.	no visible macrofauna due to sediment mobility	tidal current dominated may be a Protected wave exposure but shows an assemblage of indicator species from higher wave exposures. An assemblage observed "anomalous" for the wave energy of the site.	<i>Balanus glandula</i> <i>Fucus distichus</i>
lower	<i>Alaria</i> 'sensu morph'	<i>Hydrophyllum scabra</i>	<i>Laminaria groenlandica</i> <i>Laminaria saccharina</i> <i>Alaria</i> 'marginata' morph	<i>Laminaria saccharina</i>	<i>Laminaria groenlandica</i> <i>Laminaria saccharina</i> <i>Alaria</i> 'marginata' morph	<i>Laminaria saccharina</i>				poorly water in lagoon creates narrow intertidal and a reduced water, may have associated current dominated at outflow
subtidal	<i>Nereocystis luetkeana</i>	<i>Nereocystis luetkeana</i> <i>Macrocystis integrifolia</i> <i>Agardh</i> spp. <i>Strongylocentrotus</i> <i>Fructicosus</i>	<i>Nereocystis luetkeana</i> <i>Macrocystis integrifolia</i> <i>Agardh</i> spp. <i>Strongylocentrotus</i> <i>Fructicosus</i> <i>Zostera marina</i>	<i>Macrocystis integrifolia</i> <i>Agardh</i> spp.	<i>Nereocystis luetkeana</i> <i>Macrocystis integrifolia</i> <i>Agardh</i> spp.	<i>Macrocystis integrifolia</i> <i>Agardh</i> spp.				

