

5 - Bedrock/Gravel - CC 1-23,33 - P/VP Tidal Lagoon 11 - Bedrock or Sediment - CC 35 - VP/P/SP CC Type

Rock Shore Types - characterized by a lack of clastic sediments such as gravel or sand.

Sediment Shore Types - have substrates that have little or no bedcrock cropping out 1 Rock Ramp, Wide
2 Rock Platform Wide
3 Rock Cliff Narrow
4 Rock Ramp, Narrow
5 Rock Platform Narrow
Rock and Sediment Shore Types - rock and pockets of clastic sediments 21 Gravel Flat, Wide
22 Gravel Beach
23 Gravel Flat or Fan
24 Sand and Gravel Flat or Fan, Wide
25 Sand and Gravel Beach
26 Sand and Gravel Flat or Fan, Narrow 26 Sand and Gravel Flat or Fan, Narrow
27 Sand Beach, Wide
28 Sand Flat 6 Ramp with Gravel Beach, Wide
7 Platform with Gravel Beach, Wide
8 Cliff with Gravel Beach
9 Ramp with Gravel Beach, Narrow
10 Platform with Gravel Beach, Narrow
11 Ramp with Sand and Gravel Beach, Wide
12 Platform with Sand and Gravel Beach, Wide
13 Cliff with Sand and Gravel Beach, Wide
14 Ramp with Sand and Gravel Beach, Narrow
15 Platform with Sand and Gravel Beach, Narrow
16 Ramp with Sand Beach, Wide
17 Platform with Sand Beach, Wide
18 Cliff with Sand Beach
19 Ramp with Sand Beach, Narrow
20 Platform with Sand Beach, Narrow 6 Ramp with Gravel Beach, Wide 29 Mud Flat
30 Sand Beach, Narrow
31 Estuaries
Man-Made Materials
32 Man-made, permeable
33 Man-made, impermeable
Current Dominated
34 Channel

Each Habitat Type has typical biological features (including both an indicator species list and typical associated biobands).

To determine the Habitat Type, the biomapper looks at the along-shore Units as designated and described by the physical mapper, and 1. records the observations of the biobands in the unit and looks for indicator species,

2. □ assigns a bio-(wave) exposure category,
3. □ reviews the physical mapped information, and
4. □ assigns the Habitat Type that best describes the unit.

The Habitat Type is based on the whole unit and is similar to the physical mappers use of the 'Coastal Class' category, in that the detailed across-shore data are summarized into one attribute. The simplified category describes the features of the whole unit. Habitat Type is a summary of the biophysical classification of the whole shore unit, based on:

 □the wave exposure as indicated by the bands, and • □ the substrate types in the unit.

Legend Definitions
CC - Coastal Classification number

• □ the biobands observed,

E - Exposed - Very high wave exposure, open ocean swellsm usually fetches >500km VE - Very Exposed - Extreme high wave exposure SE - Semi Exposed - High wave exposure, open shorelines, areas between fully exposed and more sheltered, usually fetches 50 to 500km P - Protected - Low wave expsoure, sheltered inlets, usually fetches less than 10km SP - Semi Protected - Moderate wave expsoure, partly sheltered, usually fetches 10-50km

VP - Very Protected - Very low wave exposure, fethces < 1km, sheltered anchorages at heads of bays and inletes

SUBSTRATE STABILITY	IMMOBILE SUBSTRATES				MOBILE OR PARTIALLY MOBILE SUBSTRATES				CURRENT- DOMI- NATED	TIDAL IAGOON
MAJOR SUBSTRATE	BEDROCK	BEDROCK/BOULDER	BEDROCK/GRAVEL	BEDROCK/GRAVEL	SAND & GRAVEL	SAND & GRAVEL	SAND/MUD	SEDIMENT	BEDROCK OR SEDIMENT	BEDROCK OR SEDIMENT
COASTAL CLASSES	1-20	1-23, 32, 33	1-23, 33	1-23, 33	24 – 30, 32 no SAL band	24 – 30, 32 no SAL band	24 - 30, 31 has SAL band	24-30	34	35
EXPOSURE (EXP BIO)	E E	SE SE	SP	VP, P	SP SAL band	VP, P	VP, P, SP	SE, E	VP, P, SP	VP, P, SP
COMMUNITY CODE (HAB OBS)	2	3	4	5	6	7	8	9	10	11
upper middle	Verrucaria	Verrucaria Enteromorpha	Verrucaria Enteromorpha	Verrucaria Enteromorpha	Verrucaria Enteromorpha	Verrucaria Enteromorpha	grasses & rushes Salicornia virginica			
	Balanus glandula	Balanus glandula Fucus distichus	Balanus glandula Fucus distichus	Balanus glandula Fucus distichus	Balanus glandula Fucus distichus	Balanus glandula Fucus distichus	Balanus glandula Fucus distichus	no visible	tidal current	Balanus glandul Fucus distichus
	Pollicipes polymerus Mytilus californianus	Mytilus californianus	Mytilus trossulus*	Mytilus trossulus *			Mytilus trossulus*	due to sediment mobility be a Protected wave exposure but shows an assemblage of indicator species from higher wave exposures.		
	Semibalanus carriosus	Semibalanus carriosus	Semibalanus carriosus Ulva/ Ulvaria spp.	Ulva/ Ulvaria spp.	Semibalanus carriosus Ulva/ Ulvaria spp.	Ulva/ Ulvaria spp.	Ulva/ Ulvaria			ponded water in lagoon creates
mid/low	Alaria 'nana' morph	Hedophyllum sessile								narrow intertidal and a reduced biota in brackish water, may have associated current dominated at outflow
lower	Lessoniopsis littoralis  Lithothannion	Phyllospadix scouleri  Alaria 'marginata' morph Lithothamnion	Laminaria groenlandica Laminaria saccharina Alaria 'marginata' morph Lithothamnion	Laminaria saccharina	Laminaria groenlandica Laminaria saccharina Alaria 'marginala' morph Lithothamnion	Laminaria saccharina				
subtidal	Nereocystis luetkeana	Nereocystis luetkeana Macrocystis integrifolla Agarum spp. Strongylocentrotus franciscanus	Nereocystis luetkeana Macrocystis integrifolia Agarum spp. Strongylocentrotus franciscanus	Macrocystis integrifolia Agarum spp.	Nereocystis luetkeana Macrocystis integrifolia Agarum spp. Strongylocentrotus franciscanus	Macrocystis integrifolia Agarum spp.				
	l	g	Zostera marina	Zostera marina	Zostera marina	Zostera marina	Zostera marina		I	I

